

Vaccine	Timing	Eligibility	Timing
Pfizer-BioNTech	Primary series using reduced dose (10 µg/0.3 mL)	<b>Do not give - Anticipate FDA authorization and CDC recommendations by November 3</b>  Ages 5 through 11	2 doses separated by 21 days
	Primary series (30 µg/0.3 mL)	Anyone age 12+	2 doses separated by 21 days
	Third dose (30 µg/0.3 mL)	<i>Must be both</i> <ul style="list-style-type: none"> <li>Ages 12+ AND</li> <li>Moderately to severely immunocompromised<sup>1</sup></li> </ul>	28+ days after second dose
	Booster using "full dose" of Pfizer (30 µg/0.3 mL)	<i>Either</i> <ul style="list-style-type: none"> <li>65+ OR</li> <li>Both 18+ AND reside or work in a long term care facility OR</li> <li>Both 18+ AND at high risk of severe COVID<sup>2</sup> OR</li> <li>Both 18+ AND work in high-risk occupations<sup>3</sup> OR</li> <li>Both 18+ AND are BIPOC (per VT DoH) OR</li> <li>Both 18+ AND live with a BIPOC (per VT DoH)</li> </ul>	6+ months after second dose
	Heterologous booster option	Those meeting eligibility criteria for booster may opt for <ul style="list-style-type: none"> <li>"Half dose" of Moderna (50 µg) OR</li> <li>"Full dose" of Johnson &amp; Johnson (5x10<sup>10</sup> vp/ 0.5 mL)</li> </ul>	6+ months after second dose

Vaccine	Timing	Eligibility	Timing
Moderna	Primary series (100 µg/0.5 mL)	Anyone age 18+	2 doses separated by 28 days
	Third dose (100 µg/0.5 mL)	<i>Must be both</i> <ul style="list-style-type: none"> <li>Ages 18+ AND</li> <li>Moderately to severely immunocompromised<sup>1</sup></li> </ul>	28+ days after second dose
	Booster using "half dose" (50 µg/0.25 mL)	<i>Either</i> <ul style="list-style-type: none"> <li>65+ OR</li> <li>Both 18+ AND reside or work in a long term care facility OR</li> <li>Both 18+ AND at high risk of severe COVID<sup>2</sup> OR</li> <li>Both 18+ AND work in high-risk occupations<sup>3</sup> OR</li> <li>Both 18+ AND are BIPOC (per VT DoH) OR</li> <li>Both 18+ AND live with a BIPOC (per VT DoH)</li> </ul>	6+ months after second dose
	Heterologous booster option	Those meeting eligibility criteria for booster may opt for <ul style="list-style-type: none"> <li>"Full dose" of Pfizer-BioNTech (30 µg) OR</li> <li>"Full dose" of Johnson &amp; Johnson (5x10<sup>10</sup> vp/ 0.5 mL)</li> </ul>	6+ months after second dose

Vaccine	Timing	Eligibility	Timing
Johnson & Johnson (Janssen Biotech)	Primary series (5x10 <sup>10</sup> vp/ 0.5 mL)	Anyone age 18+	1 dose
	Booster using "full dose" (5x10 <sup>10</sup> vp/ 0.5 mL)	Anyone age 18+	2+ months after initial dose
	Heterologous booster option	Anyone age 18+ may opt for <ul style="list-style-type: none"> <li>"Full dose" of Pfizer-BioNTech (30 µg) OR</li> <li>"Half dose" of Moderna (50 µg)</li> </ul>	2+ months after initial dose

<sup>1</sup> CDC list of moderately to severely immunocompromised conditions
<ul style="list-style-type: none"> <li>• Been receiving active cancer treatment for tumors or cancers of the blood</li> <li>• Received an organ transplant and are taking medicine to suppress the immune system</li> <li>• Received a stem cell transplant within the last 2 years or are taking medicine to suppress the immune system</li> <li>• Moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome)</li> <li>• Advanced or untreated HIV infection</li> <li>• Active treatment with high-dose corticosteroids or other drugs that may suppress your immune response</li> </ul>
<sup>2</sup> CDC list of medical conditions placing someone at high risk of severe COVID-19
<ul style="list-style-type: none"> <li>• Cancer</li> <li>• Chronic kidney disease</li> <li>• Chronic lung diseases, including COPD (chronic obstructive pulmonary disease), asthma (moderate-to-severe), interstitial lung disease, cystic fibrosis, and pulmonary hypertension</li> <li>• Dementia or other neurological conditions</li> <li>• Diabetes (type 1 or type 2)</li> <li>• Down syndrome</li> <li>• Heart conditions (such as heart failure, coronary artery disease, cardiomyopathies or hypertension)</li> <li>• HIV infection</li> <li>• Immunocompromised state (weakened immune system)</li> <li>• Liver disease</li> <li>• Overweight and obesity</li> <li>• Pregnancy</li> <li>• Sickle cell disease or thalassemia</li> <li>• Smoking, current or former</li> <li>• Solid organ or blood stem cell transplant</li> <li>• Stroke or cerebrovascular disease, which affects blood flow to the brain</li> <li>• Substance use disorders</li> </ul>
<sup>3</sup> CDC list of professions that place individuals at high risk of exposure to potentially infectious people
<ul style="list-style-type: none"> <li>• First responders (healthcare workers, firefighters, police, congregate care staff)</li> <li>• Education staff (teachers, support staff, daycare workers)</li> <li>• Food and agriculture workers</li> <li>• Manufacturing workers</li> <li>• Corrections workers and residents</li> <li>• U.S. Postal Service workers</li> <li>• Public transit workers</li> <li>• Grocery store workers</li> <li>• Food service workers</li> <li>• Homeless shelter workers and residents</li> </ul>